

## Fire Blight

The University of Arizona • College of Agriculture • Tucson, Arizona 85721

9/98

### *Plant Disease Management: Horticultural Crops*

**MARY W. OLSEN**

*Extension Plant Pathologist  
Department of Plant Pathology*

**DEBORAH J. YOUNG**

*Extension Plant Pathologist*

### **Pathogen**

Bacterium, *Erwinia amylovora*

### **Host**

Apple, pear, pyracantha, quince and other ornamentals

### **Symptoms/signs**

If infections take place in the spring during bloom, blossom clusters wilt and shrivel, turn dark brown or black and die within a few days. During spring and summer months, infected shoots and twigs have reddish-brown streaking that turns brown in apple and brown to black in pear. Extensive branch killing is more common in pear than in apple. Infected shoots often bend at the tips and are referred to as "shepherd's crooks" in describing disease symptoms. Masses of bacteria may ooze from infected tissue as a sticky amber or brown droplets. Cankers in infected branches and trunks have a water-soaked appearance, then become sunken and dry; reddish-brown streaks may appear in the bark.

### **Environmental conditions**

Disease develops during warm temperatures and high humidity, between approximately 60° F and 85° F, and relative humidity above 60% with free moisture from dew, fog, rain or irrigation. In Arizona, these conditions are often prevalent during the monsoon season in July and August, and disease is more common during this time of the year, especially at higher elevations where both temperature and humidity are favorable for pathogen growth.

### **Disease**

Fire Blight is caused by a bacterium, *Erwinia amylovora*, that can infect many plants in the Rose family but is usually only a severe disease of apple, pear and pyracantha in Arizona. The bacterium survives in previously infected tissues in the host plant. It also has been recovered from the surface of healthy plant parts, including buds and wood. When environmental conditions are right, the bacterium multiplies in the plant and infects healthy tissue.

Bacteria enter plant tissue through natural openings or wound sites. Insects, such as bees, flies and ants, splashing rain or irrigation, and pruning tools spread the bacteria to other parts of the plant: flowers, leaves, shoots and fruits. The bacterium will continue to multiply and kill the host tissue as long as environmental conditions favor pathogen growth.

Fire Blight is more likely to show up in the high desert and mountains during the summer as a twig or stem canker disease since bloom periods in the spring are usually too dry for disease development. It may be found at lower elevations earlier if there is a high enough humidity, usually created by irrigation practices.

### ***At a Glance***

- Fireblight is a bacterial disease that affects only plants in the Rose family, particularly apple, pear and pyracantha.
- Dieback of twigs and branches, as well as shriveling of flowers, occurs in warm, moist weather; affected tissues turn very dark and twigs are crook shaped.
- Prevent disease by planting tolerant varieties; on susceptible varieties, use an antibiotic or copper spray to prevent bloom infection.
- Control by pruning infected tissue well away from infection site using a 20% bleach dip for tools between cuts.

## Prevention/control

### RESISTANT VARIETIES

#### • Apples

*Resistant:* Arkansas Black, Red Astrachan

*Moderately resistant:* Golden Delicious, Red Delicious, Granny Smith, Jonagold, Gravenstein

*Susceptible:* Gala, Rome Beauty, Jonathan, Lodi, flowering crabapple.

#### • Pears

*Resistant:* Surecrop, Kieffer

*Moderately resistant:* Seckel

*Highly susceptible:* Bosc and Bartlett; Asian pears are susceptible but not damaged as severely as Bartlett

### CLEAN PROPAGATION AND PLANTING STOCK

Locate nurseries in areas away from known infestations and use trees known to be healthy for budwood.

### PRUNING

When plants are dormant in winter, remove dead tissue by cutting at least 12 inches below infected tissue. Diseased tissue also should be removed as it appears in the spring or summer. Use a 20% household bleach solution to clean tools between each cut. Pruned plant material should be removed immediately and preferably burned.

### CHEMICAL

Chemical sprays prevent bloom infection and may require several applications. Begin treatments at budding or no later than 10% bloom. Apply every 3 to 5 days until the end of bloom. Re-spray immediately after rainfall. Spraying twigs and branches later in the season is not highly effective.

Check labels carefully for phytotoxicity. Use label rates according to chemical products.

Compound	Trade name	Comments
Antibiotics Streptomycin sulfate	Agri-Step Agri-Mycin 17	resistance to streptomycin has been found; use alternatives if resistance is suspected
Terramycin	Mycoshield Oxytetracycline Biostat	use as alternative for streptomycin
Copper sulfates	Bordeaux mixture Basicop C-O-C-S	copper can cause russetting of fruit; check label for sensitivity of varieties; acceptable for organic grown
Fixed copper	Kocide	can cause russetting of fruit; acceptable for organic grown

*Any products, services, or organizations that are mentioned, shown, or indirectly implied in this publication do not imply endorsement by The University of Arizona.*