

Barley Response to Soil Water Depletion Levels, 2000

S. H. Husman, M. J. Ottman, R. J. Wegener, and M. T. Rogers

Abstract

This research represents the first year of a project to determine when to irrigate barley based on soil water depletion levels. The purpose of this work is to establish the optimum irrigation timing based on depletion of plant available water in the soil. A field experiment was conducted at the Maricopa Agricultural Center testing irrigation of barley at 35, 50, 65, and 80% depletion of plant available water in the soil for two barley varieties, Baretta and Max. Grain yields averaged over the two varieties were 8415, 7735, 7512, and 4553 lbs/acre for the 35, 50, 65, and 80% depletion levels, respectively. The results of this study indicate irrigating at 35% soil water depletion is optimal for barley grain yield.

Introduction

Barley is an important crop in Arizona since it breaks insect and disease cycles, adds organic material to the soil due to its high straw content, and may actually increase yields of subsequent crops. Growing a crop of barley can help control winter weeds since barley is very competitive with other plants. Also, salts can be leached from the soil while growing barley during the winter when water use is low.

Barley should be irrigated when 55% of the plant available water is depleted in the root zone (Doorenbos and Pruitt, 1977). When crop water use is less than 0.10 inches/day this value should be increased by 30% (to 72% allowable depletion) and when crop water use is greater than 0.30 inches/day this value should be decreased by 30% (to 38% allowable depletion). During ripening, the maximum allowable depletion of plant available water is 90%. These guidelines are not always used in commercial practice in Arizona due to practical considerations. During the winter when water use is low and water is not needed by other crops, growers sometimes apply irrigation water to relatively wet soils to leach salts and to wet the subsoil. On the other hand, during its peak water use period, irrigation intervals often increase when water is needed to pre-irrigate cotton.

Some research on barley irrigation has been conducted in Arizona. Consumptive use curves for barley have been developed and published (Erie et al., 1968). Roth et al. (1987) studied the effects of nitrogen levels and water application amounts on barley.

Apparently, research has not been conducted in Arizona to determine when to irrigate barley based on the allowable soil water depletion levels. Therefore, the purpose of this work is to establish the optimum irrigation timing based on depletion of plant available water in the soil.

Materials and Methods

A barley irrigation study was conducted at the University of Arizona Maricopa Agricultural Center on Field 6 on a Casa Grande Sandy loam soil during the 1999-00 growing season. Two barley varieties, Baretta and Max, were planted on 23 November 99 at a rate of 135 lbs seed/acre in alternating strips. An irrigation to germinate the seed was applied on 2 December 99. The first post-emergence irrigation was applied uniformly over the entire field on 25 January 99. Subsequently, irrigations were applied when 35, 50, 65, or 80% of the plant available soil water was depleted. Irrigations were applied using the border flood method and a ditch weir was used to measure the amount of

This is a part of the University of Arizona College of Agriculture and Life Sciences 2000 Forage and Grain Report. Index at: <http://ag.arizona.edu/pubs/crops/az1185/>

water applied. The experimental design was split plot consisting of four irrigation treatments as main plots, two varieties as subplots, and four replications. The subplots comprising a single variety were 9 ft. wide and 430 ft. long. The main plots, or irrigation treatments, were 18 ft. wide and 430 ft. long.

Soil water content was measured using a Campbell Pacific 503 DR Hydroprobe. Two neutron access tubes were located in each irrigation treatment 150 ft. from the top end of the field in one variety and 150 ft. from the bottom end of the field for the other variety. Soil water content was measured using the neutron probe in the 0 to 12 inch depth increment and every 8 inches thereafter to a depth of 52 inches. Soil samples were removed on 13 January 2000 for determination of gravimetric soil water content and soil texture. The volumetric soil water content was calculated assuming bulk density values based on soil texture (USDA-SCS, 1991). The neutron probe was calibrated using the volumetric soil water content and the corresponding neutron probe readings for each depth increment. Plant available water content was calculated as the difference between soil water content at field capacity and permanent wilting point. The soil water content at permanent wilting point was determined based on its texture (USDA-SCS, 1991). Soil water content was measured with the neutron probe every 2 days until the targeted soil water depletion threshold was attained. The active root zone was expanded from the initial 0 to 12 inches when water use occurred in the next 8 inch increment since the previous irrigation. The amount of irrigation water applied was that necessary to refill the soil profile to field capacity.

The amount and timing of irrigation water and fertilizer is presented in Table 1. Fertilizer was broadcast preplant at a rate of 106 lbs N/acre as ammonium sulfate and 11-52-0 and 104 lbs P₂O₅/acre as 11-52-0. Postplant nitrogen fertilizer was applied as urea ammonium nitrate solution (32-0-0) injected into the irrigation water. The center 5 feet of each plot was harvested on 15 May 2000 with a small plot combine and grain yield was calculated. Kernel weight was determined from a 10 g sample and test weight was measured using a 1 pint container. Kernels per head was determined by weighing grain from 10 heads per plot, then dividing by kernel weight. Heads per unit area was calculated from grain yield, kernel weight, and kernels per head.

Results and Discussion

The 1999-00 barley growing season can be characterized as warm and dry compared to normal (Table 2). March was the only month where rainfall was recorded. The growing season had above average temperature except for December.

The effect of irrigation timing based on soil water depletion levels on grain yield and kernel characteristics is presented in Table 3. The optimum soil water depletion level according to grain yield was 35% for each variety and averaged over varieties. The lowest yield was obtained at 80% depletion averaged over varieties, and yield at 50 and 65% depletion was intermediate. Baretta and Max responded similarly to soil water depletion.

Irrigating at higher depletion levels decreased test weight and kernel weight. Test weight is an indirect measure of kernel size and density, and shriveled kernels can result in low test weight as these results demonstrate. More frequent irrigation resulted in more kernels per head and more heads per square foot.

In conclusion, irrigating at 35% soil water depletion was required to optimize barley grain yield in this study.

Acknowledgements

Financial support for this project was provided by the Arizona Grain Research and Promotion Council.

References

- Erie, L. J., O. F. French, and K. Harris. 1968. Consumptive use of water by crops in Arizona. Ariz. Ag. Exp. Sta. Bul. No. 169. Univ. Ariz., Tucson.
- Doorenbos, J., and W. O. Pruitt. 1977. Guidelines for predicting crop water requirements. FAO irrigation and drainage paper no. 24. Food Agric. Org. UN, Rome.
- Roth, R., B. Gardner, and B. Tickes. 1987. Barley response to water and nitrogen levels. p. 100-105. Forage and Grain. Univ. Ariz. Coll. Agric. Coop. Ext. Report Series P-71. Univ. Ariz., Tucson.
- USDA-SCS. 1991. Soil-plant-water relationships. National Engineering Handbook, Sec. 15, Ch. 1.

Table 1. Irrigation and nitrogen application dates and amounts.

Treatment	Date	Irrigation applied inches	Nitrogen applied lbs/acre
35% P.A.W. Depletion	12/2/99	4	106
	1/25/00	4	25
	2/4/00	3	0
	2/23/00	3	35
	3/21/00	3	17.5
	4/6/00	3	0
	4/18/00	4	0
	Total	24	184
50% P.A.W. Depletion	12/2/99	4	106
	1/25/00	4	25
	2/14/00	3	35
	3/21/00	4	17.5
	4/13/00	4	0
	4/18/00	4	0
	Total	23	184
65% P.A.W. Depletion	12/2/99	4	106
	1/25/00	4	25
	2/23/00	4	52.5
	3/29/00	5	0
	4/18/00	4	0
	Total	21	184
80% P.A.W. Depletion	12/2/99	4	106
	1/25/00	4	25
	3/8/00	4	70
	4/18/00	4	0
	Total	16	184

Table 2. Climatic data for Maricopa for the 1999 – 2000 growing season compared to the long-term average.

Climate variable	Year(s)	Dec	Jan	Feb	Mar	Apr	May
Max Temp. (°F)	1999-00	67	71	73	75	89	99
	Avg. ‡	67	68	71	76	84	93
Min Temp. (°F)	1999-00	32	36	39	44	52	61
	Avg. ‡	36	35	37	42	47	55
Ppt. (in)	1999-00	0.00	0.00	0.00	1.97	0.00	0.00
	Avg. ‡	1.53	0.59	0.83	0.67	0.39	0.11

‡Averages based on data summarized by Western Regional Climate Center from 1961-1990.

Table 3. Grain yield and kernel characteristics as affected by soil water depletion at irrigation (as a % of plant available water) for Max and Baretta.

Variety	Soil water depletion %	Grain yield lbs/acre	Test weight lb/bu	Kernel weight g/1000	Kernels per head	Heads per ft ²
Max	35	8435 a	54.4 a	42.8 a	41.4 ab	49.8 a
	50	7898 b	53.8 b	39.4 b	42.7 a	49.1 a
	65	7674 b	53.4 c	39.5 b	44.0 a	46.1 a
	80	4228 c	47.4 d	30.4 c	34.0 b	43.9 a
Baretta	35	8396 a	54.7 a	46.6 a	54.5 a	34.6 ab
	50	7572 b	50.1 b	42.1 b	52.3 ab	35.8 ab
	65	7350 b	50.8 b	38.2 c	51.6 ab	39.4 a
	80	4879 c	50.6 b	37.1 c	44.8 b	32.2 b
Average	35	8415 a	54.6 a	44.7 a	47.9 a	42.2 ab
	50	7735 b	51.9 b	40.8 b	47.5 a	42.4 ab
	65	7512 b	52.1 b	38.9 c	47.8 a	42.8 a
	80	4553 c	49.0 c	33.8 d	39.4 b	38.1 b

Means followed by the same letter are not significantly different according to an F-test protected LSD at P=0.05.