

Evaluation of Irrigation Termination Effects on Fiber Micronaire and Yield of Upland Cotton, 2000-2002

J.C. Silvertooth and A. Galadima

Abstract

Arizona has experienced a trend of increasing fiber micronaire values in recent years resulting in substantial discounts on fiber value. There is some evidence to suggest that irrigation termination (IT) management can influence fiber micronaire. Field studies were conducted in 2000, 2001, and 2002 at the University of Arizona Maricopa Agricultural Center (1,175ft. elevation) and in 2001 and 2002 at the Yuma Valley Agricultural Center (YVAC; 150 ft. elevation) to evaluate the effects of three dates of irrigation termination on the yield and fiber micronaire of several Upland cotton varieties. Three dates of irrigation termination (IT1, IT2, and IT3) were imposed based upon crop development into cutout. The earliest irrigation termination date, IT1 was made slightly ahead of an optimum date to provide sufficient soil-water such that bolls set at the end of the first fruiting cycle would not be water stressed and could be fully matured. Thus, the IT1 date was imposed to try to reduce overall micronaire. The second termination (IT2) date received one additional irrigation over an optimal point for the completion of the first cycle fruit set and two irrigations beyond IT1. The final (IT3) date (late September) was imposed so that soil moisture would be sufficient for the development of bolls set up through the last week of September, thus providing full top-crop potential. In general, lint yield and micronaire results revealed significant differences among the IT treatments. Micronaire and lint yield values consistently increased with later IT dates.

Introduction

One of the advantages associated with a cotton (*Gossypium spp.*) production system in an irrigated desert region such as Arizona, is the availability of a relatively long growing season, or a reliable supply of abundant heat units (HU). Traditionally, cotton production systems in the low (elevation) desert regions of Arizona (<2,000 ft. above sea level) have employed a long, full season approach. Such a long, full season approach would commonly involve a February or March date of planting with final irrigations being applied in September or October (depending on local conditions). Production over this period would include a completion of the first, or primary fruiting cycle, a cutout period (hiatus in blooming), followed by a second fruiting cycle or top-crop. Accordingly, long season, indeterminate varieties were usually best suited to this type of production system. This is one of the reasons that Pima (*G. barbadense* L.) has been well adapted to this region.

Overall, the objective with a reduced season approach to cotton production in the irrigated southwest is to achieve the highest degree of efficiency and profit to the grower as possible. To do so requires an identification of the point of diminishing returns with respect to a cotton crop. This is based on the assumption that yield potentials decline with time in the later stages of the growing season due to natural crop senescence, shorter day lengths, and cooler weather conditions (lower rates of HU accumulations).

Recent research in Arizona has attempted to address this issue by comparing a reduced season approach to that of a more traditional long, or full season system (Silvertooth et al., 1989; Silvertooth et al., 1990; Silvertooth et al., 1991; Silvertooth et al., 1992; and Silvertooth et al., 1993; Silvertooth et al., 1994; Unruh et al., 1995; Silvertooth and Norton, 1996; and Silvertooth and Norton, 1997). Summarizing this work, Unruh and Silvertooth (1997) reported on 12 site-years of data in Arizona comparing various planting and irrigation termination date combinations. The overall results from these studies revealed a most pronounced improvement in yield from an early date of planting and a generally small increase in yield from a late irrigation termination date. Comparing early and late IT treatments with an early date of planting, Unruh and Silvertooth (1997) found an average increase of 83 and 118 lbs. of lint/acre for DPL 90 and Pima S-6, respectively. Large increases in lint yield from a later IT were observed in a few experiments, but usually under conditions of very poor fruit retention over the primary fruiting cycle (up to cut-out).

About 600 HU (86/55 °F thresholds) are required to develop a late season boll from a bloom to a full sized, hard boll when fiber length development is complete (Silvertooth et al., 1996). Approximately 400 additional HU are then required to complete boll maturation and opening, for a total of 1,000 HUs needed for boll development from bloom to open boll. Therefore, IT treatments are best structured to accommodate development of bolls intended for harvest to the point of full fiber development (600 HU post-anthesis). This commonly translates to a period of approximately 21 days in southern Arizona in August and September. Accordingly, adequate soil moisture must be maintained throughout this three-week period for the last set of bolls intended for harvest. The exact IT date will therefore vary depending upon soil water holding capacities, amounts of water applied per irrigation, weather conditions, and crop condition. For example, if bolls set up to the point of cut-out are designated as those intended for harvest, final irrigations should be made so that adequate soil moisture is maintained for a three week (600 HU) period beyond the time of cut-out. The development of a top-crop usually requires irrigation and pest control for four to six weeks beyond cut-out, which for many systems equates to approximately one extra acre-foot of irrigation water and appropriate pest control to protect the developing fruit load.

In recent years an increasing percentage of the Upland (*G. hirsutum* L.) cotton crop in Arizona has been classified with micronaire (mic) ranges in excess of 4.9, resulting in a discount of the market value of the fiber. In 1999, slightly over 40% of the Arizona Upland cotton crop was classed with mic values greater than 4.9. For example, Group 6 mic values (5.0-5.2) can result in \$0.05/lb. discounts and Group 7 (≥ 5.3) \$0.10/lb. discounts. With low market values of cotton lint, as have been experienced recently (i.e. ~ \$0.50/lb. or less), discounts of this magnitude can have a devastating impact on farm revenues. Some economists have estimated that this problem has resulted in a loss of revenue to the Arizona cotton producers of approximately \$13 to 15 million per year in the past several years. However, some cotton marketing professionals in Arizona have indicated that they believe these losses in revenue due to high micronaire are in the range of \$20 to 25 million per year over the past four to five years. Thus, high micronaire is substantially reducing the profitability of Arizona cotton production in recent years.

Strategies to reduce micronaire in Arizona must consider three primary factors: 1) genetics, 2) environment, and 3) management. These three factors form a complex set of effects and interactions that determine the micronaire of a crop. The degree of genetic influence on micronaire depends on the different types of varieties that are adapted to the region in question and the range of environmental conditions within that region. The statement is often made that “only 30% of the cotton micronaire properties are determined by genetics (variety) with 70% determined by agronomic management.” University of Arizona variety trial data from 1996 through 1999 indicate that about 20% of the variation in micronaire in central Arizona was due to the genetic control of the varieties in those tests. In comparison, variation due to varieties was 50% for fiber strength, 36% for fiber length and 17% for lint yield. These data show that the environment and management exert more influence on micronaire than other fiber properties, so no matter what variety is chosen, growers can, and probably will, see wide variations in micronaire values from field to field and year to year. There remains a strong genetic component, however, the average micronaire can be reduced through proper varietal selection.

The relationship between varieties and micronaire is also complicated by the fact that high yield is genetically related to high micronaire. With our current genetic resources, the higher yielding varieties also tend to produce higher micronaire as well. This relationship between yield and micronaire is probably a

strong contributing factor to the trends we have observed in micronaire over the last few years. In Arizona, we have seen a slight increase in average mic values in the early 1990's (~1993) and again in about 1996. A similar trend is apparent with data from the entire U.S. cotton belt. Also, in review of the mic distributions among all cotton producing regions in the U.S., there is a somewhat normal distribution pattern with a peak mic value at approximately 4.9 - 5.0 and distinct drop above 5.0. These two points support the hypothesis that there is a strong genetic component associated with recent trends in Arizona and U.S. mic values and that varieties have been developed to "push" the mic limits (i.e. 5.0).

There is also ample evidence to support the position that Arizona, particularly the low elevation locations (< 2,000 ft.), has a hot environment with abundant sunlight that is conducive to high micronaire production (with hot conditions for both day and night temperatures). Thus, it appears that in Arizona we are producing a cotton crop in an environment that is conducive to high mic production with varieties that, as a whole, have a tendency toward high mic as well. The relationships associated with high mic and the third primary component (management) is not well understood in the context of desert cotton production.

Based on an analysis of data from several cotton growing areas in Arizona, it appears that there is indeed a relationship associated with location and variety and fiber mic. From this data there also appears to be a relationship between fiber mic and management, in that certain growers within given areas tend to have a very high percentage of their crop classed with high mic and another set of growers in the same area have a very low percentage of their crop with low mic using basically the same group of varieties.

There is evidence from earlier studies conducted in Arizona (Silvertooth et al., 1989; Silvertooth et al., 1990; Silvertooth et al., 1991; Silvertooth et al., 1992; and Silvertooth et al., 1993; Silvertooth et al., 1994; Unruh et al., 1995; Silvertooth and Norton, 1996; and Silvertooth and Norton, 1997) to study the effects of IT on yield and quality to suggest that IT and/or defoliation can have a significant impact on fiber mic. The initial efforts with this project revealed a significant reduction in fiber micronaire as a function of early IT management (Silvertooth et al., 2001).

It is the purpose of this research project to better delineate the contributions associated with genetics, environment, and management on fiber micronaire. The objective of this study was to further investigate the issue of IT management and the subsequent effects on the growth, development, yield, and micronaire of a group of common Upland varieties.

Materials and Methods

This study was conducted in 2000, 2001, and 2002 at the Maricopa Agricultural Center (MAC; 1,175 ft. elevation) on a Casa Grande sandy loam soil and at the Yuma Valley Agricultural Center (YVAC; 150 ft. elevation) on an Indio clay loam soil in 2001 and 2002. The experimental design was a split plot in a randomized complete block design with four replications. The main treatments consisted of three IT dates, designated as IT1, IT2, and IT3. Each main plot consisted of 12, 40-inch rows that extended the full length of the irrigation run (600 ft.). The subunits consisted of 13 Upland varieties at MAC and five varieties at YVAC. Subplots were 12, 40-inch rows wide, and 40 feet in length. The entire study areas were dry-planted and watered-up at both sites. All inputs such as fertilizer, water, and pest control were managed on an as-needed basis.

A complete set of plant measurements were collected inseason from all plots on 14-day intervals to monitor progress of growth and development of the crop. Measurements taken included: plant height, number of mainstem nodes, first fruiting branch, total number of aborted sites (positions 1 & 2), number of nodes above the top (1st position) fresh flower (NAWF), canopy closure, and number of blooms per unit area. Climatic conditions were also monitored using an Arizona Meteorological network (AZMET) stations located on both sites.

Irrigation termination treatments were imposed in relation to the crop fruiting cycle in a manner similar to that described in Figure 1. In tracking crop development, the crop was approaching cut-out, normally considered as having NAWF \leq 5, as evidenced by an average NAWF \sim 6 among all varieties. The first IT treatment (IT1) was made with the intention of terminating irrigations somewhat pre-maturely. Based upon current UA

recommendations for IT to complete a single cycle fruit set, the more optimal date of IT would have included one additional irrigation (beyond IT1). In this experiment, IT2 was structured to provide an additional (one) irrigation beyond the more optimal date. For the IT3 plots the intention was to attempt a second cycle fruit set and irrigations were continued until approximately 15 September. The IT2 treatment consistently received two additional irrigations over IT1 and IT3 received four additional irrigations over IT1 (approximately two acre-feet of additional irrigation water).

Crop management and treatments are outlined in Tables 1-10. The planting, IT, defoliation, and harvest dates are outlined in Tables 1-5. Only the center four rows of each 12-row plot were harvested.

Approximately 20 lb. seed cotton subsamples were collected from each plot at harvest. These subsamples were ginned for turnout estimates and submitted to the USDA Cotton Classing office in Phoenix, AZ for HVI analysis. All mic and lint yield data were subjected to appropriate analysis of variance procedures (Steel and Torrie, 1980 and the SAS Institute, SAS, 1990).

Results

Results reveal a consistent pattern with micronaire and lint yield responses in the IT X Variety experiments for the 2000, 2001, and 2002 seasons. The IT X Variety results for 2000, 2001, and 2002 are presented in Tables 11-47 and Figs. 1-21.

Summarizing the results for the 2000-2002 seasons, we found that fiber micronaire was significantly lower in the first IT treatment at both MAC and YVAC (less than 5.0). In 2001 at MAC, the third IT treatment (promoting a top crop development) did not produce a significant increase in yield as it did in 2000 and currently in 2002 (Figs. 5, 9, and 13). In addition, in 2001 at YVAC, the third IT treatment resulted in slightly higher but not significant increase ($p > 0.05$) in lint yield with the top-crop. In 2002 the top crop did produce a significant yield increase ($p < 0.05$) at both MAC and YVAC (Figs. 17 and 21).

These results clearly support the hypothesis concerning fiber development, micronaire, and late season irrigation management. Results have consistently shown for these three seasons that fiber micronaire can be significantly reduced and held below the discount range (5.0) for most varieties by IT management (Figs. 3, 7, 11, 15, and 19). In each case studied thus far, reasonable lint yields have still been realized with the earlier IT treatments. However, these studies have also demonstrated the potential of producing significantly higher lint yield with later IT, usually at the expense of higher fiber micronaire (≥ 5.0) values. A regression analysis of all data for the three years of this project at each location revealed a significant linear relationship for both micronaire and lint yield in relation to IT treatments (Figs. 22 and 23).

These results are consistent with earlier work on this topic (Silvertooth et al., 1989; Silvertooth et al., 1990; Silvertooth et al., 1991; Silvertooth et al., 1992; and Silvertooth et al., 1993; Silvertooth et al., 1994; Unruh et al., 1995).

Acknowledgements

The financial support provided by the Arizona Cotton Growers Association and Cotton Inc., the Arizona Cotton Research and Protection Council, Anderson Clayton Company, and Handwerker-Winburne is greatly appreciated. We gratefully acknowledge the excellent assistance from the personnel at the Maricopa and Yuma Valley Agricultural Centers. Also the hard work and technical assistance provided by the research assistants with the UA Cotton Agronomy program is greatly appreciated.

References

- SAS Institute. 1990. SAS procedure guide. Version 6, 3rd ed. SAS Inst., Cary, NC.
- Silvertooth, J.C. 2001. Recent yield and fiber micronaire tendencies for Upland cotton in Arizona. p. 41-47. Cotton, Univ. of Arizona Rep. P-125.
- Silvertooth, J.C. and A. Galadima. 2002. Evaluation of irrigation termination effects on fiber micronaire and yield of Upland cotton, 2001-2002. p. 17-41. Cotton, Univ. of Arizona Rep. P-130.
- Silvertooth, J.C., A. Galadima, E.R. Norton, and H. Moser. 2001. Evaluation of irrigation termination effects on fiber micronaire and yield of Upland cotton, 2000. p. 13-24. Cotton, Univ. of Arizona Rep. P-125.
- Silvertooth J. C., J. E. Malcuit, D. R. Howell and P. Else. 1989. Effect of date of planting on the lint yield of several cotton varieties planted at four locations in Arizona, 1988. p. 69-72. Cotton, Univ. of Arizona Rep. P-77.
- Silvertooth J. C., T. F. Watson, J. E. Malcuit, and P. W. Brown. 1992. Evaluation of date of planting and irrigation termination in the yield of Upland and Pima cotton. p. 252-273. Cotton, Univ. of Arizona Rep. P-91.
- Silvertooth J. C., T. F. Watson, J. E. Malcuit, and P. W. Brown. 1993. Evaluation of date of planting and irrigation termination in the yield of Upland and Pima cotton. p. 27-39. Cotton, Univ. of Arizona Rep. P-94.
- Silvertooth J. C., T. F. Watson, L. I. Terry, and J. E. Malcuit. 1990. Evaluation of date of planting and irrigation termination in the yield of Upland and Pima cotton. p. 6-12. Cotton, Univ. of Arizona Rep. P-81.
- Silvertooth J. C., T. F. Watson, L. I. Terry, and J. E. Malcuit. 1991. Evaluation of date of planting and irrigation termination in the yield of Upland and Pima cotton. p. 1-14. Cotton, Univ. of Arizona Rep. P-87.
- Silvertooth, J.C., and E.R. Norton. 1996. Evaluation of irrigation termination management on yield of upland cotton, 1995. p. 40-45. Cotton, Univ. of Arizona Rep. P-103.
- Silvertooth, J.C., and E.R. Norton. 1997. Evaluation of irrigation termination management on yield of upland cotton, 1996. p. 218-224. Cotton, Univ. of Arizona Rep. P-108.
- Silvertooth, J.C., E.R. Norton, and P.W. Brown. 1996. Cotton growth and development patterns. p. 75-97. Cotton, Univ. of Arizona Rep. P-103.
- Steel, R.G.D., and J.H. Torrie. 1980. Principles and procedures of statistics. McGraw-Hill, New York.
- Unruh, B.L., J.C. Silvertooth, P.W. Brown, and E.R. Norton. 1995. Effect of planting date on yield of Upland and Pima cotton varieties at Marana. p. 20-24. Cotton, Univ. of Arizona Rep. P-96.
- Unruh, B.L. and J.C. Silvertooth. 1997. Planting and irrigation termination timing effects on the yield of Upland and Pima cotton. J. Prod. Agric. 10:74-97.

Figure 1. General irrigation termination points in relation to the fruiting cycle.




Table 1. Management summary for irrigation termination by variety experiment (all plots were watered up on 6 April), MAC, 2000.

Treatment	Irrigation Termination Date	Defoliation Date	Harvest Date	Irrigation Added
1	25 July	12 August	15 September	0
2	17 August	15 September	4 October	2
3	15 September	12 October	5 December	4

Table 2. Management summary for irrigation termination by variety experiment (all plots were watered up on 17 April), MAC, 2001.

Treatment	Irrigation Termination Date	Defoliation Date	Harvest Date	Irrigation Added
1	1 August	24 August	12 September	0
2	24 August	14 September	3 October	2
3	20 September	26 October	21 November	4

Table 3. Management summary for irrigation termination by variety experiment (all plots were watered up on 4 April), MAC, 2002.

Treatment	Irrigation Termination Date	Defoliation Date	Harvest Date	Irrigation Added
1	16 July	16 August	5 September	0
2	5 August	10 September	3 October	2
3	16 September	24 October	19 November	4

Table 4. Management summary for irrigation termination by variety experiment (all plots were watered up on 21 March), YVAC, 2001.

Treatment	Irrigation Termination Date	Defoliation Date	Harvest Date	Irrigation Added
1	12 July	16 August	4 September	0
2	27 July	5 September	24 September	2
3	20 September	19 October	14 November	4

Table 5. Management summary for irrigation termination by variety experiment (all plots were watered up on 10 April), YVAC, 2002.

Treatment	Irrigation Termination Date	Defoliation Date	Harvest Date	Irrigation Added
1	18 July	30 August	13 September	0
2	15 August	4 September	13 September	2
3	15 September	10 October	25 October	4

Table 6. Irrigation termination dates and varieties, irrigation termination dates by variety study, MAC, 2000 (6 April planting and water-up).

Varieties	Maturity Type
BXN47	Medium
DP428B	Short-medium
DP422BR	Short
DP33B	Full
DP20B	Early
DP388	Medium
STV474	Medium
STV4691B	Medium
SG125BR	Short
SG747	medium
DP655BR	Full
DP451BR	medium
DP5415	Full
Irrigation Termination Dates	
Date 1 (25 July)	(3162 HU/Jan. 1)
Date 2 (17 August)	(3765 HU/Jan. 1)
Date 3 (15 September)	(4623 HU/Jan. 1)

Table 7. Irrigation termination dates and varieties, irrigation termination dates by variety study, MAC, 2001 (17 April planting and water-up).

Varieties	Maturity Type
BXN47	Medium
DP428B	Short-medium
DP422BR	Short
DP33B	Full
DP458BR	Early
DP388	Medium
STV474	Medium
STV4691B	Medium
SG215BR	Short
SG747	medium
DP655BR	Full
DP451BR	medium
DP5415	Full
Irrigation Termination Dates	
Date 1 (1 August)	(3200 HU/Jan. 1)
Date 2 (24 August)	(3843 HU/Jan. 1)
Date 3 (20 September)	(4554 HU/Jan. 1)

Table 8. Irrigation termination dates and varieties, irrigation termination dates by variety study, MAC, 2002 (2 April planting and water-up).

Varieties	Maturity Type
BXN47	Medium
DP428B	Short-medium
DP422BR	Short
DP33B	Full
DP458BR	Early
DP388	Medium
STV474	Medium
STV4691B	Medium
SG215BR	Short
SG747	medium
DP655BR	Full
DP451BR	medium
DP5415	Full
AG3601	Full
Irrigation Termination Dates	
Dates 1 (16 July)	(2795 HU/Jan. 1)
Dates 2 (5 August)	(3367 HU/Jan. 1)
Dates 3 (16 September)	(4487 HU/Jan. 1)

Table 9. Irrigation termination dates and varieties, irrigation termination dates by variety study, YVAC, AZ, 2001 (21 March planting and water-up).

Varieties	Maturity Type
SG501BR	Short
SG747	medium
STV474	medium
DP22BR	short
DP451BR	medium
Irrigation Termination Dates	
Date 1 (12 July)	(2783 HU/Jan. 1)
Date 2 (27 July)	(3195 HU/Jan. 1)
Date 3 (20 September)	(4763 HU/Jan. 1)

Table 10. Irrigation termination dates and varieties, irrigation termination dates by variety study, YVAC, AZ, 2002 (9 April planting and water-up).

Varieties	Maturity Type
SG501BR	Short
SG747	medium
STV474	medium
DP22BR	short
DP451BR	medium
Irrigation Termination Dates	
Dates 1 (18 July)	(2922 HU/Jan. 1)
Dates 2 (15 August)	(3722 HU/Jan. 1)
Dates 3 (15 September)	(4570 HU/Jan. 1)

Table 11. Experimental effects and statistical significance from the analysis of variance on micronaire, irrigation termination by variety study, MAC, 2000.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0032
Variety	0.0012
Irrigation Termination Date * Variety	0.5247

Table 12. Main effect results of micronaire for irrigation termination dates and varieties, MAC,2000.

Irrigation Termination Date	Micronaire
1	4.58 b*
2	5.23 a
3	5.18 a
LSD	0.16
OSL**	0.0032
CV(%)§	6.99
Variety	
DP655BR	5.32 a
DP5415	5.20 ab
DP422BR	5.17 ab
DP33B	5.17 ab
DP388	5.13 ab
BXN47	5.12 abc
DP451BR	5.02 abc
STV474	4.97 abc
DP428B	4.93 abcd
STV4691B	4.82 bcd
SG747	4.81 bcd
DP20B	4.73 cd
SG125BR	4.54 d
LSD	0.40
OSL	0.0012
CV(%)	8.6

*Least Significant Difference – means followed by the same letter are not significantly different according to a Fishers mean separation test at 0.05 level.

**Observed Significance Level.

§Coefficient of Variation

Table 13. Micronaire results for all varieties by irrigation termination dates, MAC, 2000.

Irrigation Termination Date 1 (24 July)	Micronaire
DP 655BR	5.07
DP 5415	4.97
DP 388	4.87
DP 33B	4.80
DP 451BR	4.77
BXN 47	4.70
DP 428B	4.53
DP 422BR	4.53
STV 4691B	4.47
STV 474	4.43
SG 125BR	4.17
SG 747	4.13
DP 20B	4.10
Irrigation Termination Date 2 (15 August)	
DP 422BR	5.53
DP 655BR	5.50
DP 5415	5.43
BXN 47	5.40
DP 33B	5.33
SG 747	5.20
DP 451BR	5.17
STV 474	5.17
DP 428B	5.13
DP 388	5.13
STV 4691B	5.13
DP 20B	5.00
SG 125BR	4.83
Irrigation Termination Date 3 (15 September)	
DP 422BR	5.43
DP 388	5.40
DP 655BR	5.40
DP 33B	5.37
STV 474	5.30
BXN 47	5.27
DP 5415	5.20
DP 428B	5.13
DP 451BR	5.13
DP 20B	5.10
SG 747	5.10
STV 4691B	4.87
SG 125BR	4.63

Irrigation Termination Date * Variety = NS (OSL = 0.5247)




Figure 2. Micronaire values as affected by irrigation termination date for each variety, MAC, 2000.




Figure 3. Mean micronaire values as affected by irrigation termination dates for all varieties, MAC, 2000.

Table 14. Micronaire results by variety for each irrigation termination date, MAC, AZ, 2000.

Irrigation Termination Date	BXN 47	DP 20B	DP 33B	DP 388	DP 422BR	DP 428B	DP 451BR	DP 5415	DP 655BR	SG 125BR	SG 747	STV 4691B	STV 474
1	4.70	4.10	4.80	4.87	4.53	4.53	4.77	4.97	5.07	4.17	4.13	4.47	4.43
2	5.27	5.00	5.33	5.13	5.53	5.13	5.17	5.43	5.50	4.83	5.20	5.13	5.17
3	5.40	5.10	5.37	5.40	5.43	5.13	5.13	5.20	5.40	4.63	5.10	4.87	5.30

Irrigation Termination Date * Variety = NS (OSL = 0.5247)

Table 15. Experimental effects and statistical significance from the analysis of variance on lint yield, irrigation termination by variety, MAC, 2000.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0272
Variety	0.0529
Irrigation Termination Date * Variety	0.0452

Table 16. Yield results (lbs. lint/acre) by variety for each irrigation termination date for cotton planted on 6 April, MAC, AZ, 2000.

Irrigation Termination Date	BXN 47	DP 20B	DP 33B	DP 388	DP 422BR	DP 428B	DP 451BR	DP 5415	DP 655BR	SG 125BR	SG 747	STV 4691B	STV 474
1	1245	1271	1131 b	1233	1283	1194 b	1359	1131	1042 b	1093 b	1385 b	1448	1334 b
2	1441	1614	1614 a	1525	1550	1617 a	1534	1527	1385 ab	1578 a	1225 b	1708	1347 b
3	1626	1741	1855 a	1531	1677	1829 a	1842	1601	1690 a	1728 a	1829 a	1855	1702 a
LSD*	NS	NS	449	NS	NS	370	NS	NS	450	423	413	NS	274
OSL**	0.2212	0.0664	0.0260	0.115	0.1466	0.0211	0.1149	0.1904	0.0402	0.0302	0.338	0.1243	0.0334
CV (%)§	15.3	11.4	12.9	4.3	12.9	10.6	13.6	19.2	14.5	127	12.3	11.1	8.3

*Least Significant Difference – means followed by the same letter within a column are not significantly different according to a Fishers mean separation test at the 0.05 level.

**Observed Significance Level.

§Coefficient of Variation.

Table 17. Yield results for all varieties by irrigation termination dates, MAC, 2000.

Irrigation Termination Date 1 (24 July)		Lint Yield (lbs. lint/acre)
	STV4691B	1448 a
	SG747	1385 ab
	DP451BR	1359 ab
	STV474	1334 abc
	DP422BR	1283 abcd
	DP20B	1270 abcd
	DP388	1245 abcde
	BXN47	1245 abcde
	DP428B	1194 bcde
	DP5415	1131 cde
	DP33B	1131 cde
	SG125BR	1093 de
	DP655BR	1042 e
LSD*		204
OSL**		0.0099
CV (%)		9.7
Irrigation Termination Date 2 (15 August)		
	STV4691B	1708
	DP428B	1619
	DP33B	1613
	DP20B	1613
	SG125BR	1578
	DP422BR	1550
	DP388	1537
	DP451BR	1535
	DP5415	1527
	BXN47	1441
	DP655BR	1385
	STV474	1347
	SG747	1225
LSD*		NS
OSL**		0.0991
CV (%)		11.1
Irrigation Termination Date 3 (15 September)		
	DP33B	1855
	ST4691B	1855
	DP451BR	1842
	DP428B	1829
	SG747	1829
	DP20B	1741
	SG125BR	1728
	ST474	1702
	DP655BR	1690
	DP422BR	1677
	BXN47	1626
	DP5415	1601
	DP388	1544
LSD*		NS
OSL**		0.2218
CV (%)§		8.5

*Least Significant Difference (LSD)

**Observed Significance Level at the 0.05 level.

§Coefficient of Variation




Figure 4. Lint yield as affected by irrigation Termination Dates for each variety, MAC, 2000.




Figure 5. Mean lint yield as affected by irrigation termination dates for all varieties, MAC, 2000.

Table 18. Experimental effects and statistical significance from the analysis of variance on micronaire, irrigation termination by variety, MAC, 2001.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0010
Variety	<0.0001
Irrigation Termination Date * Variety	0.0160

Table 19. Main effect results of micronaire for irrigation termination dates and varieties for cotton planted on 17 April (water-up) MAC, 2001.

Irrigation Termination Date	Micronaire
1	4.74 b*
2	5.07 a
3	5.19 a
LSD	0.14
OSL**	0.0160
CV(%)§	6.28
Variety	
STV474	5.33 a
SG747	5.33 a
BXN47	5.11 ab
SG215BR	5.08 ab
DP451BR	5.08 ab
DP428B	5.07 ab
STV4691B	5.00 b
DP388	5.00 b
DP33B	4.97 bc
DP458BR	4.88 bcd
DP422BR	4.84 bcd
DP5415	4.70 cd
DP655BR	4.61 d
LSD	0.40
OSL	<0.0001
CV(%)	8.6

*Least Significant Difference – means followed by the same letter are not significantly different according to a Fishers mean separation test at 0.05 level.

**Observed Significance Level.

§Coefficient of Variation

Table 20. Micronaire results for all varieties by irrigation termination dates, MAC, 2001.

Irrigation Termination Date 1 (1 August)		Micronaire
	SG 747	5.13 a*
	STV 474	5.07 a
	SG 215BR	4.97 ab
	DP 388	4.97 ab
	DP 428B	4.93 ab
	BXN 47	4.90 abc
	DP 451BR	4.86 abc
	STV 4691B	4.73 abcd
	DP 33B	4.60 bcde
	DP 422BR	4.67 cde
	DP 458BR	4.37 de
	DP 655BR	4.37 de
	DP 5415	4.23 e
LSD*		0.4567
OSL**		0.0033
CV (%)		5.72
Irrigation Termination Date 2 (24 August)		
	SG 747	5.57 a
	STV 474	5.47 ab
	BXN 47	5.30 abc
	SG 215BR	5.27 abc
	DP 451BR	5.17 abcd
	DP 33B	5.07 bcd
	STV 4691	5.03 cd
	DP 422BR	5.00 cd
	DP 428B	5.00 cd
	DP 458BR	4.93 cd
	DP 5415	4.93 cd
	DP 388	4.80 de
	DP 655BR	4.43 e
LSD*		0.4047
OSL**		0.0009
CV (%)		4.73
Irrigation Termination Date 3 (20 September)		
	STV 474	5.47 a
	DP 458BR	5.33 ab
	SG 747	5.30 abc
	DP 428B	5.27 abc
	DP 33B	5.23 abc
	STV 4691	5.23 abc
	DP 388	5.23 abc
	DP 451BR	5.20 abcd
	BXN 47	5.13 abcde
	DP 422BR	5.07 bcde
	DP 655BR	5.03 cde
	SG 215BR	5.00 de
	DP 5415	4.93 e
LSD*		0.2759
OSL**		0.0254
CV (%)§		3.12

*Least Significant Difference (LSD)

**Observed Significance Level at the 0.05 level.

§Coefficient of Variation




Figure 6. Micronaire values as affected by irrigation termination date for each variety, MAC, 2001.




Figure 7. Mean micronaire values as affected by irrigation termination dates for all varieties, MAC, 2001

Table 21. Micronaire results by variety for each irrigation termination date for cotton planted on 17 April (water-up), MAC, 2001.

Irrigation Termination Date	BXN 47	DP 33B	DP 388	DP 422BR	DP 428B	DP 451BR	DP 458BR	DP 5415	DP 655BR	SG 215BR	SG 747	STV 4691B	STV 474
1	4.90 b	4.60 b	4.97	4.47 b	4.93	4.87	4.37 c	4.23 b	4.37	4.97	5.13	4.73	5.07
2	5.30 a	5.07 a	4.80	5.00 a	5.00	5.17	4.93 b	4.93 a	4.43	5.27	5.57	5.03	5.47
3	5.13 a	5.23 a	5.23	5.07 a	5.27	5.20	5.33 a	4.93 a	5.03	5.00	5.30	5.23	5.47
LSD*	0.1999	0.4627	NS	0.1511	NS	NS	0.3294	0.5926	NS	NS	NS	NS	NS
OSL**	0.0130	0.0420	0.1008	0.0007	0.2034	0.4937	0.0032	0.0476	0.1013	0.1172	0.2139	0.878	0.1234
CV (%)§	1.73	4.11	3.65	1.38	3.86	6.81	2.98	5.56	6.66	2.86	4.66	4.00	3.90

*Least Significant Difference – means followed by the same letter within a column are not significantly different according to a Fishers mean separation test at the 0.05 level.

**Observed Significance Level.

§Coefficient of Variation.

Table 22. Experimental effects and statistical significance from the analysis of variance on lint yield, irrigation termination by variety, MAC, 2001.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0126
Variety	0.0006
Irrigation Termination Date * Variety	0.0011

Table 23. Yield results (lbs. lint/acre) by variety for each irrigation termination date for cotton planted on 17 April (water-up), MAC, 2001.

Irrigation Termination Date	BXN 47	DP 33B	DP 388	DP 422BR	DP 428B	DP 451BR	DP 458BR	DP 5415	DP 655BR	SG 215BR	SG 747	STV 4691B	STV 474
1	1302	1215 b	1258	1320	1306	1421	1156 b	953	955 b	1414	1474	1306	1271
2	1416	1625 a	1506	1581	1256	1419	1671 a	1295	1397 a	1493	1399	1467	1232
3	1211	1380 ab	1301	1385	1409	1540	1647 a	1508	1420 a	1643	1476	1503	1330
LSD*	NS	449	NS	NS	NS	NS	154	NS	389	NS	NS	NS	NS
OSL**	0.2116	0.0297	0.1012	0.1403	0.5668	0.2563	0.0012	0.0760	0.0493	0.1392	0.8697	0.3551	0.4790
CV (%)§	8.8	8.2	8.2	9.01	12.6	5.9	4.6	16.9	13.6	7.3	13.8	10.9	7.1

*Least Significant Difference – means followed by the same letter within a column are not significantly different according to a Fishers mean separation test at 0.05 level.

**Observed Significance Level.

§Coefficient of Variation.

Table 24. Yield results for all varieties by irrigation termination dates, MAC, 2001.

Irrigation Termination Date 1 (1 August)	Lint Yield (lbs. lint/acre)
SG747	1474 a*
DP451BR	1421 ab
SG215BR	1414 ab
DP422BR	1320 bc
DP428B	1306 bc
STV4691B	1306 bc
BXN47	1302 bc
STV474	1271 bcd
DP388	1258 bcd
DP458BR	1214 cd
DP33B	1156 d
DP655BR	955 e
DP5415	953 e
LSD*	150
OSL**	<0.0001
CV (%)	7.1
Irrigation Termination Date 2 (24 August)	
DP458BR	1671 a*
DP33B	1625 a
DP422BR	1581 a
DP388	1505 a
SG215BR	1493 a
STV4691B	1467 a
DP451BR	1419 a
BXN47	1415 a
SG747	1399 a
DP655BR	1397 a
DP5415	1295 a
DP428B	1256 a
STV474	1232 a
LSD*	NS
OSL**	0.0669
CV (%)	11.3
Irrigation Termination Date 3 (20 September)	
DP458BR	1647 a*
SG215BR	1643 a
DP451BR	1540 ab
DP5415	1508 abc
STV4691B	1503 abc
SG747	1476 abc
DP655BR	1420 abcd
DP428B	1409 bcd
DP422BR	1385 bcd
DP33B	1380 bcd
STV474	1330 bcd
DP388	1302 cd
BXN47	1211 d
LSD*	228
OSL**	0.0197
CV (%)§	9.4

*Least Significant Difference (LSD)

**Observed Significance Level at the 0.05 level.

§Coefficient of Variation




Figure 8. Lint yield as affected by irrigation termination dates for each variety, MAC, 2001.




Figure 9. Mean lint yield as affected by irrigation termination dates for all varieties, MAC, 2001.

Table 25. Experimental effects and statistical significance from the analysis of variance on micronaire, irrigation termination by variety, MAC, 2002.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0071
Variety	0.0014
Irrigation Termination Date * Variety	0.5648

Table 26. Main effect results of micronaire for irrigation termination dates and varieties for cotton planted on 17 April (water-up) MAC, 2002.

Irrigation Termination Date	Micronaire
1	5.14 c
2	5.36 a
3	5.26 b
LSD	0.96
OSL**	0.0071
CV(%)§	4.2
Variety	
AG3601	5.55 a
STV474	5.48 a
DP458BR	5.37 ab
DP428B	5.32 abc
SG215BR	5.31 abc
DP33B	5.29 bcd
SG747	5.24 bcde
DP5415	5.21 cde
BXN47	5.21 cde
DP451BR	5.20 cde
DP388	5.14 de
STV4691B	5.13 de
DP422BR	5.12 de
DP655BR	5.08 e
LSD	0.34
OSL	0.0014
CV(%)	4.0

*Least Significant Difference – means followed by the same letter are not significantly different according to a Fishers mean separation test at 0.05 level.

**Observed Significance Level.

§Coefficient of Variation

Table 27. Micronaire results for all varieties by irrigation termination dates, MAC, 2002.

Irrigation Termination Date 1 (16 July)	Micronaire
AG3601	5.40
STV474	5.37
DP428B	5.33
SG215BR	5.23
SG747	5.23
DP458BR	5.23
DP451BR	5.20
BXN47	5.17
DP5415	5.10
DP388	5.10
STN4691B	4.97
DP33B	4.97
DP422BR	4.90
DP655BR	4.90
Irrigation Termination Date 2 (5 August)	
AG3601	5.80
STV474	5.63
DP33B	5.57
DP458BR	5.50
DP655BR	5.37
DP428B	5.37
SG215BR	5.30
STV4691B	5.27
DP5415	5.27
DP388	5.23
DP451BR	5.23
BXN47	5.23
DP422BR	5.23
SG747	5.20
Irrigation Termination Date 3 (16 September)	
AG3601	5.45
STV474	5.43
SG215BR	5.40
DP458BR	5.37
DP33B	5.33
SG747	5.30
DP5415	5.27
DP428B	5.27
DP422BR	5.23
BXN47	5.23
DP451BR	5.17
STV4691B	5.17
DP388	5.10
DP655BR	4.97

Irrigation Termination Date * Variety = NS (OSL = 0.5648)




Figure 10. Micronaire values as affected by irrigation termination date for each variety, MAC, 2002.




Figure 11. Mean micronaire values as affected by irrigation termination dates for all varieties, MAC, 2002.

Table 28. Micronaire results by variety for each irrigation termination date, MAC, AZ, 2002.

Irrigation Termination Date	AG 3601	BXN 47	DP 33B	DP 388	DP 422BR	DP 428B	DP 451BR	DP 458BR	DP 5415	DP 655BR	SG 125BR	SG 747	STV 4691B	STV 474
1	5.40	5.17	4.97	5.10	4.90	5.33	5.20	5.23	5.10	4.90	5.23	5.23	4.97	5.37
2	5.80	5.23	5.57	5.23	5.23	5.37	5.23	5.50	5.27	5.37	5.30	5.20	5.27	5.63
3	5.45	5.23	5.33	5.10	5.23	5.27	5.17	5.37	5.27	4.97	5.40	5.30	5.17	5.43

Irrigation Termination Date * Variety = NS (OSL = 0.5648)

Table 29. Experimental effects and statistical significance from the analysis of variance on lint yield, irrigation termination by variety, MAC, 2002.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	<0.0001
Variety	0.0006
Irrigation Termination Date * Variety	0.0354

Table 30. Yield results (lbs. lint/acre) by variety for each irrigation termination date for cotton planted on 2 April, MAC, AZ, 2002.

IT Date	AG 3601	BXN 47	DP 33B	DP 388	DP 422BR	DP 428B	DP 451BR	DP 458BR	DP 5415	DP 655BR	SG 125BR	SG 747	STV 4691B	STV 474
1	873 b	1015	1020	1038	867 b	970	1018	1177	1087 b	1130 b	1071	990	1143	1031
2	1157 b	1044	1486	1160	1106 a	1003	1087	1578	1402 a	1466 a	1164	1332	1171	1040
3	1586 a	1281	1362	1238	1186 a	1230	1353	1884	1677 a	1735 a	1230	1368	1260	1209
LSD*	315	NS	NS	NS	128	NS	NS	453	310	309	NS	NS	NS	NS
OSL**	0.0204	0.1005	0.1050	0.4090	0.0051	0.1141	0.0532	0.0306	0.0157	0.0140	0.2269	0.1782	0.5895	0.1451
CV (%)§	6.1	10.9	15.8	14.4	5.37	11.6	10.29	12.9	9.8	9.4	8.0	17.7	11.4	8.8

*Least Significant Difference – means followed by the same letter within a column are not significantly different according to a Fishers mean separation test at the 0.05 level.

**Observed Significance Level.

§Coefficient of Variation.

Table 31. Yield results for all varieties by irrigation termination dates, MAC, 2002.

Irrigation Termination Date 1 (16 July)		Lint Yield (lbs. lint/acre)
DP458BR		1177
STV4691B		1143
DP655BR		1130
DP5415		1087
SG215BR		1072
DP388		1038
STV474		1031
DP33B		1020
DP451BR		1018
BXN47		1015
SG747		990
DP428B		970
AG3601		873
DP422BR		867
LSD*		NS
OSL**		0.5587
CV (%)		15.1
Irrigation Termination Date 2 (5 August)		
DP458BR		1578 a
DP33B		1486 a
DP655BR		1466 ab
DP5415		1408 abc
SG747		1332 abcd
STV4691B		1171 bcde
SG215BR		1164 bcde
DP388		1160 bcde
AG3601		1157 bcde
DP422BR		1106 cde
DP451BR		1067 cde
BXN47		1044 de
STV474		1040 de
DP428B		1003 e
LSD*		277
OSL**		0.0017
CV (%)		13.4
Irrigation Termination Date 3 (16 September)		
DP458BR		1884 a
DP655BR		1735 a
DP5415		1677 ab
AG3601		1587 abc
SG747		1368 bcd
DP33B		1362 bcd
DP451BR		1353 bcd
BXN47		1281 cd
STV4691B		1260 cd
DP388		1238 cd
DP428B		1230 cd
SG215BR		1230 cd
STV474		1209 cd
DP422BR		1186 d
LSD*		323
OSL**		0.0013
CV (%)§		13.8

*Least Significant Difference (LSD); **Observed Significance Level at the 0.05 level; §Coefficient of Variation




Figure 12. Lint yield as affected by irrigation termination dates for each variety, MAC, 2002.




Figure 13. Mean lint yield as affected by irrigation termination dates for all varieties, MAC, 2002.

Table 32. Experimental effects and statistical significance from the analysis of variance on micronaire, irrigation termination by variety, YVAC, 2001.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0190
Variety	0.0018
Irrigation Termination Date * Variety	0.3980

Table 33. Main effect results of micronaire for irrigation termination dates and varieties, for cotton planted (water-up) on 21 March YVAC, 2001

Irrigation Termination Date	Micronaire
1	4.91 b
2	5.09 a
3	4.99 a
LSD*	0.1164
OSL**	0.0190
CV(%)§	5.63
Variety	
SG 747	5.22 a
STV 474	5.18 a
SG 501BR	4.98 b
DP 451BR	4.85 bc
DP 422BR	4.77 c
LSD	0.3243
OSL	0.0018
CV(%)	4.58

*Least Significant Difference (LSD)

**Observed Significance Level at the 0.05 level.

§Coefficient of Variation

Table 34. Micronaire results for all varieties by irrigation termination dates, YVAC, 2001.

Irrigation Termination Date 1 (12 July)	Micronaire
SG 747	5.23
STV 474	5.00
DP 451BR	4.87
SG 501BR	4.75
DP 422BR	4.68
Irrigation Termination Date 2 (22 July)	
STV 474	5.35
SG 747	5.22
SG 501BR	5.13
DP 422BR	4.92
DP 451BR	4.82
Irrigation Termination Date 3 (20 August)	
STV 474	5.18
SG 747	5.15
SG 501BR	5.05
DP 451BR	4.88
DP 422BR	4.70

Irrigation Termination Date * Variety = NS (OSL = 0.3980)




Figure14. Micronaire values as affected by irrigation termination date for each variety, YVAC, 2001.




Figure 15. Mean micronaire values as affected by irrigation termination dates for all varieties, YVAC, 2001.

Table 35. Micronaire results by variety for each irrigation termination date, YVAC, 2001.

Irrigation Termination Date	DP	DP	SG	SG	STV
	422BR	451BR	501BR	747	474
	Micronaire				
1 (12 July)	4.68	4.87	4.75	5.28	5.00
2 (27 July)	4.93	4.83	5.13	5.23	5.35
3 (20 September)	4.70	4.88	5.05	5.15	5.18

Irrigation Termination Date * Variety = NS (OSL = 0.3980)

Table 36. Experimental effects and statistical significance from the analysis of variance on lint yield, irrigation termination by variety, YVAC, 2001.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.2801
Variety	0.5930
Irrigation Termination Date * Variety	0.1026

Table 37. Main effect results of yield for irrigation termination dates and varieties for cotton planted (water-up) on 21 March, YVAC, 2001.

Irrigation Termination Date	lbs. lint/acre
1 (17 July)	1816
2 (27 July)	1818
3 (20 September)	1942
LSD	NS
OSL**	0.2801
CV(%)§	10.9
Variety	
DP451BR	1843
STV474	1892
DP422BR	1801
SG747	1936
SG501BR	1890
LSD	NS
OSL	0.5930
CV(%)	11.7

*Least Significant Difference – means followed by the same letter are not significantly different according to a Fishers mean separation test at 0.05 level.

**Observed Significance Level.

§Coefficient of Variation

Table 38. Yield results by variety for each irrigation termination date for cotton planted on 21 March, YVAC, 2001.

Irrigation Termination Date	DP	DP	SG	SG	STV
	422BR	451BR	501BR	747	474
	lbs. lint/acre				
1 (17 July)	1785	1626	1888	1966	1817
2 (27 July)	1691	1858	1741	1936	1866
3 (20 September)	1928	2044	2040	1992	1908

Irrigation Termination Date * Variety = NS (OSL = 0.1026)

Table 39. Micronaire results for all varieties by irrigation termination dates, YVAC, 2001.

Irrigation Termination Date 1 (12 July)	Lint Yield (lbs. lint/acre)
SG 747	1966
SG 501BR	1888
STV 474	1817
DP 422BR	1785
DP 451BR	1626
Irrigation Termination Date 2 (22 July)	
STV 474	1936
SG 747	1866
SG 501BR	1858
DP 422BR	1741
DP 451BR	1691
Irrigation Termination Date 3 (20 August)	
STV 474	2044
SG 747	2040
SG 501BR	1992
DP 451BR	1927
DP 422BR	1908

Irrigation Termination Date * Variety = NS (OSL = 0.1026)




Figure 16. Lint yield as affected by irrigation termination dates for each variety, YVAC, 2001.




Figure 17. Mean lint yield as affected by irrigation termination dates for all varieties, YVAC, 2001.

Table 40. Experimental effects and statistical significance from the analysis of variance on micronaire, irrigation termination by variety, YVAC, 2002.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0009
Variety	0.0004
Irrigation Termination Date * Variety	0.3909

Table 41. Main effect results of micronaire for irrigation termination dates and varieties, for cotton planted (water-up) on 10 April, YVAC, 2002.

Irrigation Termination Date	Micronaire
1	4.63 b
2	5.16 a
3	5.02 a
LSD*	0.1405
OSL**	0.0009
CV(%)§	4.49
Variety	
DP422BR	4.66 b
SG501BR	4.96 a
STV474	5.00 a
DP451BR	5.02 a
SG747	5.03 a
LSD	0.2414
OSL	0.0004
CV(%)	5.97

*Least Significant Difference (LSD)

**Observed Significance Level at the 0.05 level.

§Coefficient of Variation

Table 42. Micronaire results for all varieties by irrigation termination dates, YVAC, 2002.

Irrigation Termination Date 1 (18 July)	Micronaire
DP451BR	4.83
SG747	4.70
STV474	4.65
SG501BR	4.58
DP422BR	4.38
Irrigation Termination Date 2 (15 August)	
SG747	5.35
STV474	5.20
SG 501BR	5.18
DP 451BR	5.15
DP 422BR	4.90
Irrigation Termination Date 3 (15 September)	
STV474	5.15
SG501BR	5.13
DP451BR	5.08
SG747	5.05
DP422BR	4.70

Irrigation Termination Date * Variety = NS (OSL = 0.3909)




Figure 18. Micronaire values as affected by irrigation termination date for each variety, YVAC, 2002.




Figure 19. Mean micronaire values as affected by irrigation termination dates for all varieties, YVAC, 2002.

Table 43. Micronaire results by variety for each irrigation termination date, YVAC, 2002.

Irrigation Termination Date	DP	DP	SG	SG	STV
	422BR	451BR	501BR	747	474
	Micronaire				
1 (18 July)	4.38	4.83	4.58	4.70	4.65
2 (15 August)	4.90	5.15	5.18	5.35	5.20
3 (15 September)	4.70	5.08	5.13	5.05	5.15

Irrigation Termination Date * Variety = NS (OSL = 0.3909)

Table 44. Experimental effects and statistical significance from the analysis of variance on lint yield, irrigation termination by variety, YVAC, 2002.

Source of Variation (Effect)	OSL (Pr >F)
Irrigation Termination Date	0.0017
Variety	0.2235
Irrigation Termination Date * Variety	0.1311

Table 45. Main effect results of yield for irrigation termination dates and varieties for cotton planted (water-up) on 10 April, YVAC, 2002.

Irrigation Termination Date	lbs. lint/acre
1	1128 b
2	1587 a
3	1607 a
LSD	
OSL**	0.0017
CV(%)§	
Variety	
DP451BR	1519
STV474	1515
SG747	1487
DP422BR	1367
SG501BR	1317
LSD	NS
OSL	0.2235
CV(%)	22.7

*Least Significant Difference – means followed by the same letter are not significantly different according to a Fishers mean separation test at 0.05 level.

**Observed Significance Level.

§Coefficient of Variation

Table 46. Yield results by variety for each irrigation termination date for cotton planted on 9 April, YVAC, 2002.

Irrigation Termination Date	DP	DP	SG	SG	STV
	422BR	451BR	501BR	747	474
	lbs. lint/acre				
1 (18 July)	1098	1260	1040	1028	1212
2 (15 August)	1408	1805	1335	1691	1695
3 (15 September)	1594	1492	1574	1741	1638

Irrigation Termination Date * Variety = NS (OSL = 0.1311)

Table 47. Micronaire results for all varieties by irrigation termination dates, YVAC, 2002.

Irrigation Termination Date 1 (18 July)	Lint Yield (lbs. lint/acre)
DP451BR	1260
STV474	1212
DP422BR	1098
SG501BR	1040
SG747	1028
Irrigation Termination Date 2 (15 August)	
DP451BR	1805
STV474	1695
SG747	1691
DP422BR	1408
SG501BR	1335
Irrigation Termination Date 3 (15 September)	
SG747	1741
STV474	1638
DP422BR	1594
SG501BR	1574
DP451BR	1492

Irrigation Termination Date * Variety = NS (OSL = 0.1311)




Figure 20. Lint yield as affected by irrigation termination dates for each variety, YVAC, 2002.




Figure 21. Mean lint yield as affected by irrigation termination dates for all varieties, YVAC, 2002.




Figure 22. Grand mean of micronaire values as affected by irrigation termination dates for all varieties, MAC, 2000-2002.




Figure 23. Grand mean of lint yields as affected irrigation termination dates for all varieties, MAC, 2000-2002.




Figure 24. Grand mean of micronaire values as affected by irrigation termination dates for all varieties, YVAC, 2001-2002.




Figure 25. Grand mean of lint yields as affected by irrigation termination dates for all varieties, YVAC, 2001-2002.




Figure 26. Relationship between fiber micronaire and heat units accumulated from planting (PD) to final irrigation (IT), MAC and YVAC, 2000-2002.




Figure 27. Relationship between lint yield and heat units accumulated from planting (PD) to final irrigation (IT), MAC and YVAC, 2000-2002.